Outline

- Sample error, true error
- Confidence intervals for observed hypothesis error
- Binomial distribution, Normal distribution, Central Limit Theorem

Two Definitions of Error

The **true error** of hypothesis h with respect to target function f and distribution $\mathcal{D}$ is the probability that h will misclassify an instance drawn at random according to $\mathcal{D}$.

$$error_{\mathcal{D}}(h) \equiv \Pr_{x \in \mathcal{D}}[f(x) \neq h(x)]$$

The **sample error** of h with respect to target function f and data sample S is the proportion of examples h misclassifies

$$error_S(h) \equiv \frac{1}{n} \sum_{x \in S} \delta(f(x) \neq h(x))$$

Where $\delta(f(x) \neq h(x))$ is 1 if $f(x) \neq h(x)$, and 0 otherwise.

How well does $error_S(h)$ estimate $error_{\mathcal{D}}(h)$?

Problems Estimating Error

---

1. *Bias:* If S is training set, $error_S(h)$ is optimistically biased

$$bias \equiv E[error_S(h)] - error_{\mathcal{D}}(h)$$

   For unbiased estimate, h and S must be chosen independently

2. *Variance:* Even with unbiased S, $error_S(h)$ may still *vary* from $error_{\mathcal{D}}(h)$

Example

Hypothesis h misclassifies 12 of the 40 examples in S

$$error_S(h) = \frac{12}{40} = .30$$

What is $error_{\mathcal{D}}(h)$?

Estimators

Experiment:

1. choose sample S of size n according to distribution $\mathcal{D}$
2. measure $error_S(h)$

$error_S(h)$ is a random variable (i.e., result of an experiment)

$error_S(h)$ is an unbiased *estimator* for $error_{\mathcal{D}}(h)$

Given observed $error_S(h)$ what can we conclude about $error_{\mathcal{D}}(h)$?

Confidence Intervals

If

- S contains n examples, drawn independently of h and each other
- $n \geq 30$

Then

- With approximately 95% probability, $error_{\mathcal{D}}(h)$ lies in interval

$$error_S(h) \pm 1.96\sqrt{\frac{error_S(h)(1 - error_S(h))}{n}}$$

Confidence Intervals

If

- S contains n examples, drawn independently of h and each other
- $n \geq 30$

Then

- With approximately N% probability, $error_{\mathcal{D}}(h)$ lies in interval

  $$error_S(h) \pm z_N \sqrt{\frac{error_S(h)(1 - error_S(h))}{n}}$$

  where

$N\%$:	50%	68%	80%	90%	95%	98%	99%
z_N:	0.67	1.00	1.28	1.64	1.96	2.33	2.58

$error_S(h)$ is a Random Variable

Rerun the experiment with different randomly drawn S (of size n)

Probability of observing r misclassified examples:



$$P(r) = \frac{n!}{r!(n-r)!}\, error_{\mathcal{D}}(h)^r (1 - error_{\mathcal{D}}(h))^{n-r}$$

Binomial Probability Distribution



$$P(r) = \frac{n!}{r!(n-r)!}\, p^r (1-p)^{n-r}$$

Probability $P(r)$ of r heads in n coin flips, if $p = \Pr(heads)$

- Expected, or mean value of X, $E[X]$, is
$$E[X] \equiv \sum_{i=0}^{n} iP(i) = np$$
- Variance of X is
$$Var(X) \equiv E[(X - E[X])^2] = np(1-p)$$
- Standard deviation of X, σ_X, is
$$\sigma_X \equiv \sqrt{E[(X - E[X])^2]} = \sqrt{np(1-p)}$$

Normal Probability Distribution



$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$

The probability that X will fall into the interval (a, b) is given by

$$\int_a^b p(x)dx$$

- Expected, or mean value of X, $E[X]$, is

$$E[X] = \mu$$

- Variance of X is

$$Var(X) = \sigma^2$$

- Standard deviation of X, σ_X, is

$$\sigma_X = \sigma$$

Normal Probability Distribution



80% of area (probability) lies in $\mu \pm 1.28\sigma$

N% of area (probability) lies in $\mu \pm z_N\sigma$

Central Limit Theorem

Consider a set of independent, identically distributed random variables $Y_1 \ldots Y_n$, all governed by an arbitrary probability distribution with mean μ and finite variance σ^2. Define the sample mean,

$$\bar{Y} \equiv \frac{1}{n} \sum_{i=1}^{n} Y_i$$

Central Limit Theorem. As $n \to \infty$, the distribution governing $\bar{Y}$ approaches a Normal distribution, with mean μ and variance $\frac{\sigma^2}{n}$.